arXiv:0801.3879v1 [gr-qc] 25 Jan 2008

Separation of Dirac equation in the $3+1$ dimensional constant curvature black hole background and its solution

Banibrata Mukhopadhyay*
Department of Physics, Indian Institute of Science, Bangalore-560012, India

Kaushik Ghosh
Department of Engineering Sciences, Haldia Institute of Technology,
Haldia, East Midnapore-721 657, West Bengal, India

The behavior of spin-half particles is discussed in the $3+1$-dimensional constant curvature black hole (CCBH) spacetime. We use Schwarzschild-like coordinates, valid outside the black hole event horizon. The constant time surfaces corresponding to the time-like Killing vector are degenerate at the black hole event horizon and also along an axis. We write down the Dirac equation in this spacetime using Newman-Penrose formalism which is not easily separable unlike that in the Kerr metric. However, with a particular choice of basis system the equation is separable and we obtain the solutions. We discuss the structural difference in the Dirac equation in the CCBH spacetime with that in the Kerr geometry, due to difference in the corresponding spacetime metric, resulting complexity arised in separation in the earlier case.

PACS numbers: 04.62.+v, 04.70.-s, 04.70.Dy, 95.30.Sf
Keywords: constant curvature black hole spacetime, spin-1/2 particle, Dirac equation, separability

I. INTRODUCTION

A new type of black hole solution has been found by Bañados [1]. This solution results from an identification of spacetime points in the anti-de Sitter space and represents a higher dimensional generalization of the $2+1$ dimensional BTZ black hole [2]. This is a constant curvature black hole (CCBH) with a negative cosmological constant. The corresponding spacetime geometry is $R^3 \times S^1$. Unlike the BTZ black hole, the $3+1$ dimensional black hole does not have rotating solution and is dynamic. However Schwarzschild-like coordinates, valid outside the horizon, can be found which covers a sub-manifold of the space time.

The scalar field solution in the $3+1$ dimensional CCBH was discussed in [3] while studying their thermodynamic behavior. On the other hand, the spinor field solution was studied extensively in black hole spacetimes over last three decades by various authors including one of the present ones (e.g. [4, 5, 6, 7, 8, 9, 10, 11, 12]). These authors discussed in detail the separability of the Dirac equation in black hole spacetimes and solutions. In the present article we study the behavior of a spin-half field in the $3+1$ dimensional CCBH spacetime. We use Newman-Penrose formalism to establish the Dirac equation. It is found that in this background the Dirac equation is not easily separable, unlike the situation in the Kerr geometry, unless we change the basis appropriately. This is, as we discuss in the following sections in detail, due to very nature of the background geometry which is, by definition, quite different from conventional black hole spacetimes.

The paper is organized as follows. In the next section we recall the $3+1$ CCBH metric and describe its properties briefly. In §III we calculate the basis vectors of null tetrad in this spacetime and write down the Dirac equation following the Newman-Penrose formalism. We show that the set of equations is separable into radial and angular part by defining new spinors obtained with a suitable combination of original spinors. Subsequently we discuss the solutions in §IV. The section V is devoted in obtaining the effective potentials of spinors in the CCBH spacetime and comparing with that in the Kerr geometry. We show that due to difference in structure of spacetime metric the Dirac equation in the CCBH spacetime appears different than that in the Kerr geometry hindering its separation in regular basis where particles are expressible as *pure* up and/or down spinors. Finally, we summarize our results in §VI.

---

*Electronic address: bm@physics.iisc.ernet.in

II. DESCRIPTION OF THE $3+1$ CCBH METRIC

The anti-de Sitter spacetime in 3+1 dimension is defined as the universal covering space of the hypersurface

$$-x_0^2 + x_1^2 + x_2^2 + x_3^2 - x_4^2 = -l^2, \tag{1}$$

where $l^2 = -1/\Lambda$, when Λ is the cosmological constant. The 3+1 dimensional CCBH is obtained by making identifications in this space using an one dimensional subgroup of its isometry group SO(2,3). In the Kruskal coordinates the metric for the 3+1 dimensional CCBH is given by [1],

$$ds^2 = \frac{l^2(r+r_h)^2}{{r_h}^2} dy^\alpha dy^\beta \eta_{\alpha\beta} + r^2 d\phi^2 \tag{2}$$

where r is given by

$$r = r_h \frac{(1+y^2)}{(1-y^2)} \tag{3}$$

and $y^2 = y^\alpha y^\beta \eta_{\alpha\beta}$ [$\eta_{\alpha\beta} =$ diag(-1,1,1)], r_h is the black hole horizon. The coordinate ranges are $-\infty < y^\alpha < \infty$ and $0 \leq \phi < 2\pi$.

We introduce the local Schwarzschild-like coordinates (t, r, θ) in the 2+1 dimensional hyperplane y^α as

$$\begin{aligned} y^0 &= \alpha(r)\cos\theta \sinh(\frac{r_h t}{l}) \\ y^1 &= \alpha(r)\cos\theta \cosh(\frac{r_h t}{l}) \\ y^2 &= \alpha(r)\sin\theta \end{aligned} \tag{4}$$

where $\alpha(r) = [\frac{(r-r_h)}{(r+r_h)}]^{\frac{1}{2}}$.

In these coordinates the metric becomes

$$ds^2 = \frac{l^4 f^2(r)}{r_h^2}[d\theta^2 - \cos^2\theta (dt/l)^2] + \frac{dr^2}{f^2(r)} + r^2 d\phi^2 \tag{5}$$

where $f(r) = (\frac{r^2-r_h^2}{l^2})^{\frac{1}{2}}$. These coordinates are valid outside the horizon $(r > r_h)$ for $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$ and $0 \leq \phi < 2\pi$. However in these coordinates only part of the space is covered $(-1 < y_2 < 1)$. It is clear that the foliation becomes degenerate along the direction $\theta = -\frac{\pi}{2}$ and $\theta = \frac{\pi}{2}$.

III. DIRAC EQUATION IN THE CCBH METRIC

In this section we will use the Newman-Penrose formalism [13] to write the Dirac equation in the above mentioned constant curvature black hole spacetime. The Dirac equation in Newman-Penrose formalism can be written as [5]

$$\sigma^\mu_{AB'} D_\mu P^A + i\mu_p \bar{Q}^{C'} \epsilon_{C'B'} = 0, \tag{6}$$

$$\sigma^\mu_{AB'} D_\mu Q^A + i\mu_p \bar{P}^{C'} \epsilon_{C'B'} = 0, \tag{7}$$

where, for any vector X_i, according to the spinor formalism $\sigma^i_{AB'} X_i = X_{AB'}$, $\epsilon_{C'B'}$ is the twodimensional Levi-Civita tensor; $A, B = 0, 1$ and

$$D_\mu P^A = \partial\mu P^A + \Gamma^A_{\mu\nu} P^\nu. \tag{8}$$

Here we introduce a null tetrad $(\vec{l}, \vec{n}, \vec{m}, \vec{\bar{m}})$ to satisfy orthogonality relations, $\vec{l}.\vec{n} = 1$, $\vec{m}.\vec{\bar{m}} = -1$ and $\vec{l}.\vec{m} = \vec{n}.\vec{m} = \vec{l}.\vec{\bar{m}} = \vec{n}.\vec{\bar{m}} = 0$ following Newman & Penrose [13]. $2^{\frac{1}{2}}\mu_p$ is the mass of the Dirac particle. In terms of this new basis, in Newman-Penrose formalism, the Pauli matrices can be written as [7]

$$\sigma^{\mu}_{AB'} = \frac{1}{\sqrt{2}} \begin{pmatrix} l^{\mu} & m^{\mu} \\ \bar{m}^{\mu} & n^{\mu} \end{pmatrix}. \tag{9}$$

Now following previous works [7, 12], writing various spin coefficients by their named symbols [7], and choosing

$$P^0 = F_1, P^1 = F_2, \bar{Q}^{1'} = G_1, \bar{Q}^0 = -G_2$$

we obtain

$$l^{\mu}\partial_{\mu}F_1 + \bar{m}^{\mu}\partial_{\mu}F_2 + (\epsilon - \tilde{\rho})F_1 + (\pi - \alpha)F_2 = i\mu_p G_1, \tag{10}$$

$$m^{\mu}\partial_{\mu}F_1 + n^{\mu}\partial_{\mu}F_2 + (\mu - \gamma)F_2 + (\beta - \tau)F_1 = i\mu_p G_2, \tag{11}$$

$$l^{\mu}\partial_{\mu}G_2 - m^{\mu}\partial_{\mu}G_1 + (\epsilon^* - \tilde{\rho}^*)G_2 - (\pi^* - \alpha^*)G_1 = i\mu_p F_2, \tag{12}$$

$$n^{\mu}\partial_{\mu}G_1 - \bar{m}^{\mu}\partial_{\mu}G_2 + (\mu^* - \gamma^*)G_1 - (\beta^* - \tau^*)G_2 = i\mu_p F_1. \tag{13}$$

These are the Dirac equations in Newman-Penrose formalism in curved space-time in absence of electromagnetic interaction. The corresponding equations in presence of electromagnetic interaction can be found in earlier works (e.g. [11, 14, 15]).

Now we calculate the basis vectors of null tetrad in terms of elements of the CCBH metric given as

$$l^{\mu} = \left(\frac{r_h}{l\cos\theta f^2}, 1, 0, 0\right), \tag{14a}$$

$$n^{\mu} = \left(\frac{r_h}{2l\cos\theta}, -\frac{f^2}{2}, 0, 0\right), \tag{14b}$$

$$m^{\mu} = \left(0, 0, \frac{r_h}{\sqrt{2}l^2 f}, \frac{i}{\sqrt{2}r}\right), \tag{14c}$$

$$\bar{m}^{\mu} = \left(0, 0, \frac{r_h}{\sqrt{2}l^2 f}, -\frac{i}{\sqrt{2}r}\right) \tag{14d}$$

and

$$l_{\mu} = \left(\frac{l\cos\theta}{r_h}, -\frac{1}{f^2}, 0, 0\right), \tag{15a}$$

$$n_{\mu} = \left(\frac{f^2 l\cos\theta}{2r_h}, \frac{1}{2}, 0, 0\right), \tag{15b}$$

$$m_{\mu} = \left(0, 0, -\frac{fl^2}{\sqrt{2}r_h}, -\frac{ir}{\sqrt{2}}\right), \tag{15c}$$

$$\bar{m}_{\mu} = \left(0, 0, -\frac{fl^2}{\sqrt{2}r_h}, \frac{ir}{\sqrt{2}}\right). \tag{15d}$$

We also calculate the various spin coefficients as

$$\tilde{\rho} = -\frac{1}{2}\left(\frac{\partial_r f}{f} + \frac{1}{r}\right), \qquad \beta = 0, \qquad \pi = -\frac{r_h \tan\theta}{2\sqrt{2}l^2 f}, \qquad \epsilon = 0,$$

$$\tau = \frac{r_h \tan\theta}{2\sqrt{2}l^2 f}, \qquad \mu = -\frac{f^2}{4}\left(\frac{\partial_r f}{f} + \frac{1}{r}\right), \qquad \gamma = \frac{f\partial_r f}{2}, \qquad \alpha = 0. \tag{16}$$

Now we consider the spin-$\frac{1}{2}$ wave function as the form of $e^{i(\sigma t + m\phi)}F(r,\theta)$, where σ is the frequency of the incoming wave and m is the azimuthal quantum number.

Thus using eqns. (14), (15), and (16), eqns. (10) to (13) reduce to

$$\left[\frac{i\sigma r_h}{l\cos\theta f} + f\partial_r + \frac{1}{2}\left(\partial_r f + \frac{f}{r}\right)\right]F_1 + \left[\frac{r_h}{\sqrt{2}l^2}\partial_\theta - \frac{r_h\tan\theta}{2\sqrt{2}l^2} + \frac{mf}{\sqrt{2}r}\right]F_2 = i\mu_p f G_1, \tag{17a}$$

$$\left[\frac{ir_h\sigma}{2l\cos\theta} - \frac{f^2}{2}\partial_r - \frac{f^2}{4}\left(\frac{\partial_r f}{f} + \frac{1}{r}\right) - \frac{f\partial_r f}{2}\right]F_2 + \frac{1}{f}\left[\frac{r_h}{\sqrt{2}l^2}\partial_\theta - \frac{r_h\tan\theta}{2\sqrt{2}l^2} - \frac{mf}{\sqrt{2}r}\right]F_1 = i\mu_p G_2, \tag{17b}$$

$$\left[\frac{i\sigma r_h}{l\cos\theta f} + f\partial_r + \frac{1}{2}\left(\partial_r f + \frac{f}{r}\right)\right]G_2 - \left[\frac{r_h}{\sqrt{2}l^2}\partial_\theta - \frac{r_h\tan\theta}{2\sqrt{2}l^2} - \frac{mf}{\sqrt{2}r}\right]G_1 = i\mu_p f F_2, \tag{17c}$$

$$\left[\frac{ir_h\sigma}{2l\cos\theta} - \frac{f^2}{2}\partial_r - \frac{f^2}{4}\left(\frac{\partial_r f}{f} + \frac{1}{r}\right) - \frac{f\partial_r f}{2}\right]G_1 - \frac{1}{f}\left[\frac{r_h}{\sqrt{2}l^2}\partial_\theta - \frac{r_h\tan\theta}{2\sqrt{2}l^2} + \frac{mf}{\sqrt{2}r}\right]G_2 = i\mu_p F_1. \tag{17d}$$

Now we define

$$\frac{\sqrt{2}F_1}{f} = \tilde{f}_1, F_2 = f_2, G_1 = g_1, \text{ and } \frac{\sqrt{2}G_2}{f} = -\tilde{g}_2, \tag{18}$$

and

$$\mathcal{D} = f^2\left(\partial_r + \frac{1}{2r}\right) + \frac{3}{2}f\partial_r f, \;\; \mathcal{L} = \frac{r_h}{l^2}\partial_\theta - \frac{r_h\tan\theta}{2l^2}, \tag{19}$$

and reduce eqns. (17a-d) to

$$\left(\mathcal{D} + \frac{i\sigma r_h}{l\cos\theta}\right)\tilde{f}_1 + \left(\mathcal{L} + \frac{mf}{r}\right)f_2 = i\sqrt{2}\mu_p f g_1, \tag{20a}$$

$$\left(\mathcal{D} - \frac{i\sigma r_h}{l\cos\theta}\right)f_2 - \left(\mathcal{L} - \frac{mf}{r}\right)\tilde{f}_1 = i\sqrt{2}\mu_p f \tilde{g}_2, \tag{20b}$$

$$\left(\mathcal{D} + \frac{i\sigma r_h}{l\cos\theta}\right)\tilde{g}_2 + \left(\mathcal{L} - \frac{mf}{r}\right)g_1 = -i\sqrt{2}\mu_p f f_2, \tag{20c}$$

$$\left(\mathcal{D} - \frac{i\sigma r_h}{l\cos\theta}\right)g_1 - \left(\mathcal{L} + \frac{mf}{r}\right)\tilde{g}_2 = -i\sqrt{2}\mu_p f \tilde{f}_1. \tag{20d}$$

We now follow the Chandrasekhar's procedure [5, 7] which was generalised by Page [6], Carter and McLenaghan [16]. The last two authors showed explicitly that the method consists of two steps. First, one has to replace the original wave equation by a modified but equivalent one such that the operator gets split up into radial and angular part which commute each other. Second, making use of the explicit form of the modified operators, one has to factorize the components of the spinor into pairs of single variable, one of r and other of θ, interdependent functions obeying a system of ordinary differential equations involving a separation constant λ. Hence, the radial and angular operators also commute with the total wave operator transformed from the original one. This implies the existence of a conserved current associated with each solution of the equation.

Carter and McLenaghan [16] showed, with the use of Penrose-Floyd tensor [17], that occurrence of type of operators mentioned above arises in presence of an appropriate Killing spinor field in the spacetime under consideration. The existence of a Killing spinor, K_{CD}, was demonstrated earlier [18] in any type-D vacuum spacetime of a second order symmetric two-spinor satisfying

$$\nabla_{A'(B}K_{CD)} = 0, \tag{21}$$

when the latin indices run from 0 to 1, parenthesis indicates symmetrization over the indices within, and the primed spinors transform under the conjugate of the transformation from the unprimed ones in a Lorentz transformation. The solution of above equation is a conformal Killing spinor, since it describes a constant motion along the null geodesic. On the other hand, constants of all the Kinnersley type-D vacuum solutions can be derived from separability of the null geodesic Hamilton-Jacobi equation [16]. It is therefore appropriate to demand the symmetric two-spinor to satisfy

$$\nabla_{A'C}K^{C}_{B} + \bar{\nabla}_{BC'}\bar{K}^{C'}_{A'} = 0. \tag{22}$$

This later condition was understood more clearly by the Penrose-Floyd tensor (see [16] and [17] for details),

$$f_{\mu\nu} \leftrightarrow \bar{\epsilon}_{A'B'}K_{AB} + \epsilon_{AB}\bar{K}_{A'B'}, \tag{23}$$

satisfying

$$f_{\mu(\nu;\delta)} = 0. \tag{24}$$

We find that defining $\Psi_1 = \tilde{f}_1 + f_2$, $\Psi_2 = f_2 - \tilde{f}_1$, $\Psi_3 = g_1 + \tilde{g}_2$ and $\Psi_4 = g_1 - \tilde{g}_2$ and combining eqns. (20a-d) we obtain

$$\left(\mathcal{D} + \frac{mf}{r}\right)\Psi_1 + \left(\mathcal{L} - \frac{i\sigma r_h}{l\cos\theta}\right)\Psi_2 = i\sqrt{2}\mu_p f\Psi_3, \tag{25a}$$

$$\left(\mathcal{D} - \frac{mf}{r}\right)\Psi_2 - \left(\mathcal{L} + \frac{i\sigma r_h}{l\cos\theta}\right)\Psi_1 = -i\sqrt{2}\mu_p f\Psi_4, \tag{25b}$$

$$\left(\mathcal{D} - \frac{mf}{r}\right)\Psi_3 + \left(\mathcal{L} - \frac{i\sigma r_h}{l\cos\theta}\right)\Psi_4 = -i\sqrt{2}\mu_p f\Psi_1, \tag{25c}$$

$$\left(\mathcal{D} + \frac{mf}{r}\right)\Psi_4 - \left(\mathcal{L} + \frac{i\sigma r_h}{l\cos\theta}\right)\Psi_3 = i\sqrt{2}\mu_p f\Psi_2 \tag{25d}$$

what satisfy all the above conditions to separate them into radial and angular operators allowing to follow the procedure by Chandrasekhar [7] generalized by Carter and McLenaghan [16] described above.

Now defining $\Psi_1 = R_-(r)S_-(\theta)$, $\Psi_2 = R_+(r)S_+(\theta)$, $\Psi_3 = R_+(r)S_-(\theta)$ and $\Psi_4 = R_-(r)S_+(\theta)$ we separate the set of equations into radial and angular parts given by

$$\left(\mathcal{D} + \frac{mf}{r}\right)R_- - i\sqrt{2}\mu_p f R_+ = \lambda_1 R_+; \quad \left(\mathcal{L} - \frac{i\sigma r_h}{l\cos\theta}\right)S_+ = -\lambda_1 S_-, \tag{26a}$$

$$\left(\mathcal{D} - \frac{mf}{r}\right)R_+ + i\sqrt{2}\mu_p f R_- = \lambda_2 R_-; \quad \left(\mathcal{L} + \frac{i\sigma r_h}{l\cos\theta}\right)S_- = \lambda_2 S_+, \tag{26b}$$

$$\left(\mathcal{D} - \frac{mf}{r}\right)R_+ + i\sqrt{2}\mu_p f R_- = \lambda_3 R_-; \quad \left(\mathcal{L} - \frac{i\sigma r_h}{l\cos\theta}\right)S_+ = -\lambda_3 S_-, \tag{26c}$$

$$\left(\mathcal{D} + \frac{mf}{r}\right)R_- - i\sqrt{2}\mu_p f R_+ = \lambda_4 R_+; \quad \left(\mathcal{L} + \frac{i\sigma r_h}{l\cos\theta}\right)S_- = \lambda_4 S_+. \tag{26d}$$

From the radial part of eqns. (26a-d) it is clear that for unique solution $\lambda_4 = \lambda_1$ and $\lambda_3 = \lambda_2$. Thus further eqns. (26a-d) reduce to

$$\left(\mathcal{D} + \frac{mf}{r}\right) R_- = \left(\lambda_1 + i\sqrt{2}\mu_p f\right) R_+; \quad \left(\mathcal{D} - \frac{mf}{r}\right) R_+ = \left(\lambda_2 - i\sqrt{2}\mu_p f\right) R_-, \tag{27a}$$

$$\left(\mathcal{D} - \frac{mf}{r}\right) R_+ = \left(\lambda_2 - i\sqrt{2}\mu_p f\right) R_-; \quad \left(\mathcal{D} + \frac{mf}{r}\right) R_- = \left(\lambda_1 + i\sqrt{2}\mu_p f\right) R_+, \tag{27b}$$

$$\left(\mathcal{L} - \frac{i\sigma r_h}{l\cos\theta}\right) S_+ = -\lambda_1 S_-; \quad \left(\mathcal{L} + \frac{i\sigma r_h}{l\cos\theta}\right) S_- = \lambda_2 S_+, \tag{27c}$$

$$\left(\mathcal{L} - \frac{i\sigma r_h}{l\cos\theta}\right) S_+ = -\lambda_2 S_-; \quad \left(\mathcal{L} + \frac{i\sigma r_h}{l\cos\theta}\right) S_- = \lambda_1 S_+. \tag{27d}$$

If we choose $\lambda_1 = \lambda_2 = \lambda$ and rename

$$\mathcal{D} + \frac{mf}{r} \rightarrow \mathcal{D}^\dagger; \quad \mathcal{D} - \frac{mf}{r} \rightarrow \mathcal{D} \tag{28a}$$

and

$$\mathcal{L} + \frac{i\sigma r_h}{l\cos\theta} \rightarrow \mathcal{L}^\dagger; \quad \mathcal{L} - \frac{i\sigma r_h}{l\cos\theta} \rightarrow \mathcal{L}, \tag{28b}$$

then eqns. (27a-d) reduce to

$$\mathcal{D}R_+ = \left(\lambda - i\sqrt{2}\mu_p f\right) R_-; \quad \mathcal{D}^\dagger R_- = \left(\lambda + i\sqrt{2}\mu_p f\right) R_+, \tag{29}$$

$$\mathcal{L}S_+ = -\lambda S_-; \quad \mathcal{L}^\dagger S_- = \lambda S_+. \tag{30}$$

The eqns. (29-30) are the separated set of Dirac equation, where $\mathcal{D}^\dagger$, $\mathcal{D}$ and $\mathcal{L}^\dagger$, $\mathcal{L}$ act as creation and annihilation operators for radial and angular functions respectively. For comparison, one may check the corresponding equations in the Kerr geometry [7, 10] given below in eqns. (47) and (48).

IV. SOLUTION FOR THE ANGULAR AND RADIAL DIRAC EQUATION

The angular equation for S_+ is given by

$$\partial_\theta{}^2 S_+ - \tan\theta\, \partial_\theta S_+ + \left[\nu(\nu+1) - \frac{n^2 - 2n\mu\sin\theta + \mu^2}{\cos^2\theta}\right] S_+ = 0 \tag{31}$$

where $n = 1/2$, $\mu = -i\sigma l$ and $\nu(\nu+1) = \frac{l^4\lambda^2}{{r_h}^2} - 1/4$. Following previous works [19, 20], the solution of eqn. (31) is given by

$$S_+(x) \rightarrow \left(\frac{1-x}{2}\right)^{\frac{n-\mu}{2}} \left(\frac{1+x}{2}\right)^{\frac{n+\mu}{2}} F(a, b, c; \frac{1-x}{2}) \tag{32}$$

where

$$a = -\nu + \frac{1}{2}[(n-\mu) + (n+\mu)],$$

$$b = \nu + 1 + \frac{1}{2}[(n-\mu) + (n+\mu)],$$

$$c = (n-\mu)+1,$$

$$x = \cos\theta. \tag{33}$$

This solution is well-behaved throughout the interval $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$ [21], the range of θ where the coordinate system is valid outside of the black hole. The solution for S_- is similar to S_+ with μ replaced by $-\mu$.

The radial equation for R_+ is given by,

$$f^2(r)\partial_r(f^2\partial_r R_+) + f^2(r)\partial_r([g(r)-h(r)]R_+) + f^2(r)[g(r)+h(r)]\partial_r R_+ + [g^2(r)-h^2(r)]R_+$$

$$-(\lambda^2 + 2\mu_p^2 f^2)R_+ + \frac{i\sqrt{2}\mu_p f^2 \partial_r f}{\lambda - i\sqrt{2}\mu_p f}(f^2\partial_r + g - h)R_+ = 0 \tag{34}$$

where $g(r) = \frac{f^2(r)}{2r} + \frac{3}{2}f(r)\partial_r f(r)$ and $h(r) = \frac{mf(r)}{r}$. The wave function $R_-(r)$ satisfies an equation similar to that of $R_+(r)$ with $h(r)$ replaced by $-h(r)$.

For $r \to \infty$, the radial equations for R_+ and R_- both reduce to

$$[\partial_r^2 + \frac{5}{r}\partial_r + \frac{4}{r^2} - (\lambda^2 + 2\mu_p^2 f^2)\frac{l^4}{r^4}]R_\pm = 0, \tag{35}$$

keeping leading order terms which gives

$$\left(\partial_r^2 + \frac{5}{r}\partial_r + \frac{4-2\mu_p^2 l^2}{r^2}\right)R_\pm = 0. \tag{36}$$

Choosing $z = -\frac{1}{4r^4}$, this reduces to

$$\left(z^2\partial_z^2 + \frac{2-\mu_p^2 l^2}{8}\right)R_\pm = 0 \tag{37}$$

whose solution is

$$R_\pm \to z^{1/2+s}, z^{1/2-s}, \tag{38}$$

where $s = \frac{\mu_p l}{2\sqrt{2}}$. Therefore, the general solutions are $R_+ = Az^{1/2+s} + Bz^{1/2-s}$ and $R_- = Cz^{1/2+s} + Dz^{1/2-s}$ when A, B, C, D are arbitrary constants. Hence for $r \to \infty$, $|R_+|$ and $|R_-|$ (as well as $|R_+|^2$ and $|R_-|^2$) both decay to zero provided $\mu_p l < \sqrt{2}$. It also follows from eqns. (36-38) that asymptotically the first derivatives of R_+ and R_- should reduce to zero and thus the asymptotic solutions for R_+ and R_- are same with $A = C$ and $B = D$ [23]. Now once we know $S_\pm$ and $R_\pm$, we can obtain original components of spinor by combining them as

$$\tilde{f}_1 = \frac{S_-R_- - S_+R_+}{2} \to r^{-2\mp 4s} F\left(a,b,c;\frac{1-x}{2}\right)\left[\left(\frac{1-x}{2}\right)^{\frac{n+\mu}{2}}\left(\frac{1+x}{2}\right)^{\frac{n-\mu}{2}}\right.$$

$$\left. - \left(\frac{1-x}{2}\right)^{\frac{n-\mu}{2}}\left(\frac{1+x}{2}\right)^{\frac{n+\mu}{2}}\right] \tag{39}$$

for $r \to \infty$. Similarly, other components f_2, g_1, $\tilde{g}_2$ can be obtained.

V. EFFECTIVE POTENTIALS OF $R_\pm$ FIELDS AND COMPARISON WITH POTENTIALS OF SIMILAR FIELDS IN THE KERR GEOMETRY

A. Effective potentials of $R_\pm$

Let us define

$$f^2 dy = dr$$

such that the eqn. (34) reduces to

$$\left[\partial_y^2+\left(2g+\frac{i\sqrt{2}\mu_p(\partial_y f)}{\lambda-i\sqrt{2}\mu_p f}\right)\partial_y+\left(g^2-h^2+\partial_y(g-h)-(\lambda^2+2\mu_p^2f^2)+\frac{i\sqrt{2}\mu_p(\partial_y f)}{\lambda-i\sqrt{2}\mu_p f}(g-h)\right)\right]R_+=0. \tag{40}$$

Defining further

$$u=\int\frac{\lambda-i\sqrt{2}\mu_p f}{rf^3}dy=\frac{\lambda l^5}{3r_h^5}\left(\frac{3r^2r_h-4r_h^3}{l^3f^3}-3\tan^{-1}\left(\frac{r_h}{lf}\right)\right)+\frac{i\sqrt{2}\mu_p l^4}{2r_h^4}\left(\frac{r_h^2}{l^2f^2}+2\log\left(\frac{lf}{r}\right)\right) \tag{41}$$

eqn. (40) reduces to

$$\left(\frac{\partial^2}{\partial u^2}+V_+\right)R_+=0 \tag{42}$$

where the effective potential

$$V_+=\left(g^2-h^2+\partial_y(g-h)-(\lambda^2+2\mu_p^2f^2)+\frac{i\sqrt{2}\mu_p(\partial_y f)}{\lambda-i\sqrt{2}\mu_p f}(g-h)\right)\frac{r^2f^6}{(\lambda-i\sqrt{2}\mu_p f)^2}. \tag{43}$$

Similarly the potential V_- for R_- can be obtained which is same as V_+ except $h(r)$ replaced by $-h(r)$. Figure 1 describes behavior of V_+ in the complex plane of u for two values of cosmological constant $\Lambda=-1/l^2$. In Fig. 2, we show V_+ for massless fermion. It is found that close to the black hole horizon when $Re(u)\to-\infty$ (and $r\to r_h$) the potential barrier vanishes, while at far away it diverges when $Re(u)\to 0$ (and $r\to\infty$). This assures that the radial solution vanishes asymptotically what we indeed show in eqns. (37)-(39). Behavior of V_- is very similar to V_+. Note that for all the figures we choose the separation constant $\lambda=1$. In principle λ should have computed explicitly, especially from the angular Dirac equation, which might have been different than unity. However, this does not affect the qualitative feature of $V_\pm$ and thus we keep the exact computation of λ as a future mission.

B. Effective potentials in the Kerr geometry

First of all let us recall the Dirac equation in Kerr geometry [7]

$$\mathcal{D}_0 f_1+2^{-1/2}\mathcal{L}_{1/2}f_2=(i\mu_* r+a\mu_*\cos\theta)g_1, \tag{44a}$$

$$\Delta\mathcal{D}^\dagger_{1/2}f_2-2^{1/2}\mathcal{L}^\dagger_{1/2}f_1=-2(i\mu_* r+a\mu_*\cos\theta)g_2, \tag{44b}$$

$$\mathcal{D}_0 g_2-2^{-1/2}\mathcal{L}^\dagger_{1/2}g_1=(i\mu_* r-a\mu_*\cos\theta)f_2, \tag{44c}$$

$$\Delta\mathcal{D}^\dagger_{1/2}g_1+2^{1/2}\mathcal{L}_{1/2}g_2=-2(i\mu_* r-a\mu_*\cos\theta)f_1, \tag{44d}$$

where f_1,f_2,g_1,g_2 are components of spinor and

$$\mathcal{D}_n=\partial_r+\frac{iK}{\Delta}+2n\frac{r-M}{\Delta}, \tag{45a}$$

$$\mathcal{D}^\dagger_n=\partial_r-\frac{iK}{\Delta}+2n\frac{r-M}{\Delta}, \tag{45b}$$

$$\mathcal{L}_n=\partial_\theta+Q+n\mathrm{cot}\theta, \tag{46a}$$

$$\mathcal{L}^\dagger_n=\partial_\theta-Q+n\mathrm{cot}\theta, \tag{46b}$$



FIG. 1: Variation of effective potential, (a) real part of V_+, (b) imaginary part of V_+, in the complex plane of u for $l = 3$ (solid curve), 1 (dashed curve). Other parameters are $M = 1$, $r_h = l$, $m = 1/2$, $\mu_p = 1$.



FIG. 2: Variation of V_+ for massless fermion as a function of (a) u, (b) r, for $l = 3$ (solid curve), 1 (dashed curve). Other parameters are $M = 1$, $r_h = l$, $m = 1/2$.

with $K = (r^2 + a^2)\sigma + am$, $Q = a\sigma\sin\theta + m\mathrm{cosec}\theta$. We follow the usual convention for various notations (see e.g. [7]) which we do not repeat here. If one compares this set of equations with the corresponding eqns. (20a-d) obtained for the CCBH spacetime, then it is easy to understand that there are differences in structure of equations. While in the CCBH background the angular and radial functions are coupled with respectively radial and angular operators, in the Kerr metric they are completely decoupled due to very nature of the spacetime. Therefore, after choosing $f_1(r,\theta) = R_{-1/2}(r)S_{-1/2}(\theta)$, $f_2(r,\theta) = R_{1/2}(r)S_{1/2}(\theta)$, $g_1(r,\theta) = R_{1/2}(r)S_{-1/2}(\theta)$, $g_2(r,\theta) = R_{-1/2}(r)S_{1/2}(\theta)$, the

Dirac equation in Kerr geometry can easily be separated into radial and angular part given by [5, 7, 10]

$$\Delta^{\frac{1}{2}}\mathcal{D}_0 R_{-\frac{1}{2}} = (\lambdabar + i m_p r)\Delta^{\frac{1}{2}} R_{+\frac{1}{2}}, \tag{47a}$$

$$\Delta^{\frac{1}{2}}\mathcal{D}_0^{\dagger}\Delta^{\frac{1}{2}} R_{+\frac{1}{2}} = (\lambdabar - i m_p r) R_{-\frac{1}{2}}, \tag{47b}$$

$$\mathcal{L}_{\frac{1}{2}} S_{+\frac{1}{2}} = -(\lambdabar - a m_p \cos\theta) S_{-\frac{1}{2}}, \tag{48a}$$

$$\mathcal{L}_{\frac{1}{2}}^{\dagger} S_{-\frac{1}{2}} = +(\lambdabar + a m_p \cos\theta) S_{+\frac{1}{2}}, \tag{48b}$$

where $m_p = 2^{1/2}\mu_*$ is the mass of the particle, $\lambdabar$ is the separation constant and $2^{1/2}R_{-1/2}$ is redefined as $R_{-1/2}$.

However, this is not obvious in the CCBH geometry. In this case, we change the basis by combining the basic spinors and obtain eqns. (25a-d) which are *mathematically* analogous to the set of eqns. (44a-d). Now as Ψ-s in eqns. (25a-d) are linear combination of spinors rather than the spinors itself, we are not able to separate the equations for radial and angular spinors, $R_{\pm 1/2}(r)$ and $S_{\pm 1/2}(\theta)$, as could do in the Kerr spacetime. However, we obtain the separated radial and angular equations for spinors in new basis given by eqns. (29)-(30).

In the Kerr geometry we change the independent variable r to r_* such that

$$r_* = r + \frac{2Mr_+ + am/\sigma}{r_+ - r_-}\log\left(\frac{r}{r_+} - 1\right) - \frac{2Mr_- + am/\sigma}{r_+ - r_-}\log\left(\frac{r}{r_-} - 1\right) \tag{49}$$

and choose $R_{-\frac{1}{2}} = P_{-\frac{1}{2}}$, $\Delta^{\frac{1}{2}} R_{+\frac{1}{2}} = P_{+\frac{1}{2}}$. Then by defining

$$(\lambdabar \pm i m_p r) = exp(\pm i\theta)\sqrt{\lambdabar^2 + m_p^2 r^2},\ \tan\theta = \frac{m_p r}{\lambdabar}, \tag{51}$$

$$P_{\pm\frac{1}{2}} = \psi_{\pm\frac{1}{2}}\ exp\left[\mp\frac{1}{2} i\ \tan^{-1}\left(\frac{m_p r}{\lambdabar}\right)\right], \tag{50}$$

$$Z_{\pm} = \psi_{+\frac{1}{2}} \pm \psi_{-\frac{1}{2}} \tag{52}$$

and combining the eqns. (47a-b) we obtain [7, 10, 22],

$$\left(\frac{d}{d\hat{r}_*} - W\right) Z_+ = i\sigma Z_-, \tag{53a}$$

and

$$\left(\frac{d}{d\hat{r}_*} + W\right) Z_- = i\sigma Z_+, \tag{53b}$$

where,

$$\hat{r}_* = r_* + \frac{1}{2\sigma}\tan^{-1}\left(\frac{m_p r}{\lambdabar}\right), \tag{54}$$

$$W = \frac{\Delta^{\frac{1}{2}}(\lambdabar^2 + m_p^2 r^2)^{3/2}}{\omega^2(\lambdabar^2 + m_p^2 r^2) + \lambdabar m_p \Delta/2\sigma},\ \ \omega^2 = \frac{K}{\sigma}. \tag{55}$$

Physically the radial eqns. (53a-b) are similar to the set of eqns. (29) obtained for the CCBH metric, except the independent variable $\hat{r}_*$ is Cartesian like while r is the radius vector of polar coordinate system. Either set is describing the spinor fields not in the original basis but in the transformed one like the linear combination of original ones. Decoupling eqns. (53a-b) we obtain

$$\left(\frac{d^2}{d\hat{r}_*{}^2} + \sigma^2\right) Z_{\pm} = V_{k\pm} Z_{\pm}, \tag{56}$$

where

$$V_{k\pm} = W^2 \pm \frac{dW}{d\hat{r}_*}$$

$$= \frac{\Delta^{\frac{1}{2}}(\lambdabar^2 + m_p^2 r^2)^{3/2}}{[\omega^2(\lambdabar^2 + m_p^2 r^2) + \lambdabar m_p \Delta/2\sigma]^2}[\Delta^{\frac{1}{2}}(\lambdabar^2 + m_p^2 r^2)^{3/2} \pm ((r - M)(\lambdabar^2 + m_p^2 r^2) + 3m_p^2 r\Delta)]$$

$$\mp \frac{\Delta^{\frac{3}{2}}(\lambdabar^2 + m_p^2 r^2)^{5/2}}{[\omega^2(\lambdabar^2 + m_p^2 r^2) + \lambdabar m_p \Delta/2\sigma]^3}[2r(\lambdabar^2 + m_p^2 r^2) + 2m_p^2 \omega^2 r + \lambdabar m_p (r - M)/\sigma] \tag{57}$$

which carries similar information as eqn. (42) does for R_+. Now if we compare eqns. (53a-b), which generate V_{k+} in eqn. (57), with eqns. (29) generating V_+ in eqn. (42), then the difference is obvious. Hence, the effective potential $V_\pm$ for $R_\pm$ fields in the CCBH spacetime come out to be different (which are complex functions) than $V_{k\pm}$ for $Z_\pm$ fields in the Kerr spacetime. Note that unlike V_+ described in Figs. 1, 2, $V_{k\pm}$ attain a finite value at $\hat{r}_* \to -\infty$ (close to the black hole horizon) and ∞ [7, 10]. In addition, by the very nature of spacetime and corresponding construction one can obtain the original spinors $R_{\pm 1/2}$ from $Z_\pm$. However, there is no obvious way to obtain original spinors in the CCBH spacetime from $R_\pm$, what one needs to define for the sake of separation, which is understood from eqns. (20), (25), (28) as described before.

VI. SUMMARY

We have considered the spin-half fermion field in the 3 + 1-dimensional CCBH background. The Dirac equation is obtained by using the Newman-Penrose formalism. We show that the Dirac equation is separable only with the change of basis. The new spinors are linear combination of original spinors giving rise to the Dirac equation separable into radial and angular parts. We obtain the corresponding solutions. We discuss the difference in the Dirac equation in the CCBH spacetime with that in the Kerr geometry recalling the geometrical conditions to be satisfied to separate them out [16]. We show that the structure of equations in the earlier case is different than the later one (due to difference in structure of the spacetimes) leading to a different separation and solution procedure. Now one can consider the Dirac equation solution in the CCBH spacetime to study the thermodynamical properties of spinor field as was done earlier for scalar field [3].

Note that the coordinate system used in the text does not cover the entire manifold outside the horizon completely. There are other foliations which cover the outside horizon region completely [24]. However in this foliation the metric becomes explicitly time dependent.

---

[1] M. Bañados, Phys. Rev. D **57**, 1068 (1998)
[2] M. Bañados, M. Henneaux, C. Teiltelboim and J. Zanelli, Phys. Rev. D **48**, 1506 (1993)
[3] K. Ghosh, Phys. Rev. D **60**, 104003 (1999)
[4] S. A. Teukolsky, Phys. Rev. Lett. **29**, 1114 (1972).
[5] S. Chandrasekhar, Proc. R. Soc. Lond. A **349**, 1 (1976).
[6] D. N. Page, Phys. Rev. **D 14**, 1509 (1976).
[7] S. Chandrasekhar, in *The Mathematical Theory Of Black Holes* (London: Clarendon Press, 1983).
[8] J. Guven, and D. Nunez, Phys. Rev. **D 42**, 2577 (1990).
[9] B. Mukhopadhyay, and S. K. Chakrabarti, Class. Quantum Grav. **16**, 3165 (1999).
[10] B. Mukhopadhyay, and S. K. Chakrabarti, Nucl. Phys. B **582**, 627 (2000).
[11] B. Mukhopadhyay, and N. Dadhich, Class. Quantum Grav. **21**, 3621 (2004).
[12] B. Mukhopadhyay, Class. Quantum Grav. **17**, 2017 (2000).
[13] E. Newman, and R. Penrose, J. Math. Phys. **3**, 566 (1962).
[14] R. Penrose, and W. Rindler, in *Spinors and Space-Time* (Cambridge University Prss, 1986).
[15] D. Häfner, and J.-P. Nicolas, Rev. Math. Phys. **16**, 29 (2004).
[16] B. Carter, and R. G. McLenaghan, Phys. Rev. **D 19**, 1093 (1979).
[17] R. Penrose, Ann. N. Y. Acad. Sci. **224**, 125 (1973).
[18] M. Walker, and R. Penrose, Commun. Math. Phys. **18**, 265 (1970).
[19] I. I. Cotaescu and M. Visinescu, Mod. Phys. Lett. A **15**, 145, (2000).

[20] K. Ghosh, Phys. Rev. D **67**, 124027, (2003).
[21] I. S. Gradshteyn and I. M. Ryzhik, Tables of Integrals Series and Products (Academic, New York, 1965).
[22] B. Mukhopadhyay, Ind. J. Phys. **B73**, 855 (1999); gr-qc/9910018.
[23] S. J. Avis, C. J. Isham, and D. Storey, Phys. Rev. **D 18**, 3565 (1978).
[24] S. Holst and P. Peldan, Class. Quantum Grav. **14**, 3433 (1997).